



Short Communication

In vivo horizontal dissemination of the *bla*_{KPC-2} gene carried on diverse genetic platforms among clinical isolates of Enterobacteriaceae



M.S. Anchordoqui^a, D. De Belder^{a,e}, C. Lucero^a, M. Rapoport^a, D. Faccone^{a,e}, A. Rodriguez^b, A. Di Martino^c, F. Martino^a, I. Herrero^d, F. Pasteran^a, A. Corso^a, S.A. Gomez^{a,e,*}

^a Servicio Antimicrobianos, National Reference Laboratory in Antimicrobial Resistance, Instituto Nacional de Enfermedades Infecciosas ANLIS 'Dr Carlos G. Malbrán', Buenos Aires, Argentina

^b Clínica y Maternidad Suizo Argentina, Buenos Aires, Argentina

^c Sanatorio Mitre, Buenos Aires, Argentina

^d Hospital Municipal de Urgencias, Córdoba, Argentina

^e Research Career, CONICET (Consejo Nacional de Investigaciones Científicas y Tecnológicas)

ARTICLE INFO

Article history:

Received 4 November 2014

Received in revised form 15 April 2015

Accepted 6 May 2015

Keywords:

KPC-2
Enterobacteriaceae
Tn4401a
Horizontal dissemination

ABSTRACT

This study investigated the molecular characteristics of six *bla*_{KPC}-positive Enterobacteriaceae recovered from three patients in Argentina. Antimicrobial susceptibility testing was performed following Clinical and Laboratory Standards Institute (CLSI) 2014 recommendations. Molecular characterisation of the isolates was performed by biparental conjugation, PCR, sequencing, S1 nuclease restriction, and Southern blot hybridisation with a *bla*_{KPC} probe using standard protocols and conditions. The isolates studied were as follows. Case 1: *Escherichia coli* (ECO-P1) and *Klebsiella pneumoniae* (KPN-P1) isolated from a rectal swab harboured *bla*_{KPC-2} in transposon Tn4401a on non-typeable and non-conjugative plasmids. Case 2: *Enterobacter cloacae* (ECL-P2) and *K. pneumoniae* (KPN-P2) were isolated from two blood cultures. *bla*_{KPC-2} was found in a novel genetic variant of ISKpn8-*bla*_{KPC-2}-ISKpn6-like on conjugative plasmids of IncL/M type. Case 3, *Citrobacter freundii* (CFR-P3) and *Klebsiella oxytoca* (KOX-P3) were isolated from skin and skin-structure infection. The *bla*_{KPC} gene was detected on ISKpn8- Δ *bla*_{TEM}-*bla*_{KPC-2}-ISKpn6-like located on an IncA/C conjugative plasmid. CFR-P3 and KOX-P3 harboured *bla*_{PER-2} in addition to the *bla*_{KPC} gene. In conclusion, we document the horizontal dissemination of *bla*_{KPC-2} from diverse Enterobacteriaceae clinical isolates with different genetic backgrounds. This is the first report of *E. coli* harbouring *bla*_{KPC} associated with Tn4401a in Argentina.

© 2015 International Society for Chemotherapy of Infection and Cancer. Published by Elsevier Ltd. All rights reserved.

1. Introduction

Genes encoding KPC enzymes are predominantly identified among *Klebsiella pneumoniae* isolates and increasingly among other Enterobacteriaceae and non-fermenters. The global predominance of KPC-producing bacteria has been mainly associated with the successful and hyperepidemic clone of *K. pneumoniae* sequence type (ST) 258 and single and double-locus variants such as ST11 [1]. *bla*_{KPC} genes are typically transposon-encoded (Tn4401 and its variants) and therefore have the potential to disseminate between plasmids and across bacterial species [2]. The spread of *bla*_{KPC} by horizontal genetic transfer has been previously documented in a

simultaneous blood infection [3], in colonised patients possessing KPC-3, Tn4401a and pKpQIL-IT elements [4] and through the transmission of a promiscuous plasmid carrying the KPC gene in Tn4401a [5].

In Argentina, health institutions throughout the country refer clinical strains for characterisation to the National Reference Laboratory in Antimicrobial Resistance (LNRAR) following local guidelines. In this context, six Enterobacteriaceae isolates recovered from three patients with phenotypes indicative of KPC production were sent to LNRAR for confirmation and molecular characterisation.

2. Methods

2.1. Patients and clinical isolates

The epidemiological data and clinical characteristics of the patients are shown in Table 1. Patients 1, 2 and 3 were hospitalised

* Corresponding author. Present address: Servicio Antimicrobianos, Departamento de Bacteriología, Instituto Nacional de Enfermedades Infecciosas ANLIS 'Dr Carlos G. Malbrán', Av. Vélez Sarsfield 563 (C1282AFF), Ciudad Autónoma de Buenos Aires, Argentina. Tel.: +54 11 4303 2812; fax: +54 11 4303 2812.

E-mail address: sgomez@anlis.gov.ar (S.A. Gomez).

Table 1
Epidemiological and clinical characteristics of patients infected or colonised with *bla*_{KPC}-positive isolates.

| Patient | Isolate | Isolation date | Age (years)/sex | Hospital (city) | Reason for admission | Underlying conditions | KPC-positive specimen | LOS (days) | Antibiotic treatment (during hospital stay) | Outcome |
|---------|------------------|--------------------------------|-----------------|---|----------------------|---|-----------------------|------------|--|---------|
| 1 | ECO-P1 KPN-P1 | 24 Oct. 2010 25 Oct. 2010 | 94/F | Clínica y Maternidad Suizo Argentina (Buenos Aires) | Colostomy | Colon cancer | Rectal swab | 45 | UKN | UKN |
| 2 | ECL-P2 KPN-P2 | 14 March 2011 21 March 2011 | 87/M | Sanatorio Dupuytren (Buenos Aires) | UKN | Neurological patient with MV | Blood | >12 | COL, VAN ^a | UKN |
| 3 | CFR-P3 KOX-P3 | 4 April 2012 | 52/M | Hospital Municipal de Urgencias (Córdoba) | Wound | DM, DVT, CD, bypass, supracondylar amputation | Wound | 26 | FLZ ^a , CAZ, VAN ^a , CLI ^a , CIP, SXT+TZP | Died |

LOS, length of stay; ECO, *Escherichia coli*; KPN, *Klebsiella pneumoniae*; ECL, *Enterobacter cloacae*; CFR, *Citrobacter freundii*; KOX, *Klebsiella oxytoca*; UKN, unknown; MV, mechanical ventilation; DM, diabetes mellitus; DVT, deep vein thrombosis; CD, cardiovascular disease; COL, colistin; VAN, vancomycin; FLZ, fluconazole; CAZ, ceftazidime; CLI, clindamycin; CIP, ciprofloxacin; SXT, trimethoprim/sulfamethoxazole; TZP, piperacillin/tazobactam.

^a Chemotherapy used to treat fungal infections or Gram-positive bacteria.

in 2010, 2011 and 2012, respectively, in different institutions in two cities in Argentina. The isolates belonged to five species, including *Escherichia coli* ($n = 1$), *K. pneumoniae* ($n = 2$), *Enterobacter cloacae* ($n = 1$), *Klebsiella oxytoca* ($n = 1$) and *Citrobacter freundii* ($n = 1$) (Table 1).

2.2. Susceptibility testing and detection of resistance genes

Minimum inhibitory concentrations (MICs) to imipenem, meropenem, ertapenem, cefepime, cefotaxime, ceftazidime, gentamicin, amikacin, ciprofloxacin, minocycline, colistin, fosfomycin and tigecycline were determined by agar dilution using standard methods according to Clinical and Laboratory Standards Institute (CLSI) criteria [6]. MIC interpretation was according to CLSI guidelines (Table 2A in CLSI document M100-S24 [7]) for all antimicrobials except for fosfomycin (endovenous), colistin and tigecycline, which were interpreted following European Committee on Antimicrobial Susceptibility Testing (EUCAST) recommendations. KPC production was detected phenotypically considering imipenem inhibition zones ≤ 22 mm and positive synergy between carbapenem disks and 3-aminophenyl-boronic acid (APB) and negative synergy between carbapenem disks and ethylene diamine tetra-acetic acid/sodium mercaptoacetic acid (EDTA/SMA) [8,9]. Extended-spectrum β -lactamase (ESBL) production was suspected phenotypically by positive synergy with clavulanic acid. Locally prevalent ESBL genes were confirmed by specific PCRs using standard conditions (CTX-MU1, 5'-ATGTGC AGYACCAGTAARGT-3'; CTX-MU2, 5'-TGGGTRARTARGTSACCAGA-3'; PER-2F, 5'-GTAG-TATCAGCCCAATCCCC-3'; PER-2R, 5'-CCAATAAAGGCCGTCCAT CA-3'). All resistance genes and genetic contexts were sequenced using the above primers as well as others previously reported using BigDyeTM Terminator methodology [10].

2.3. Conjugation experiments

Transfer of KPC-harboring elements was tested by biparental conjugation with *E. coli* J53 (sodium azide-resistant) on Luria-Bertani (LB) agar plates in a 1:3 donor/recipient ratio. Transconjugants were selected on LB agar plates with sodium azide (150 μ g/mL) and ampicillin (50 μ g/mL) [10].

2.4. Genetic typing

For pulsed-field gel electrophoresis (PFGE), total DNA was digested with *Xba*I and the fragments were separated using 2.2 s and 54.2 s as initial and final pulse times over 20 h. The running

was performed using a CHEF-DR[®] III Apparatus (Bio-Rad, Hercules, CA) and the gel was recorded and DNA patterns were analysed according to the criteria of Tenover et al. [10,11].

2.5. Plasmid analysis

Plasmid extraction was performed using a QIAGEN Midi Kit (QIAGEN, Hilden, Germany). Plasmids were classified by PCR-based replicon typing (PBRT) and S1 nuclease digestion of total DNA [12,13] and were analysed by Southern blot and hybridisation of the PFGE gels using a *bla*_{KPC-2}-specific probe (PCR DIG Probe Labeling Mix; Roche Applied Science, Barcelona, Spain).

2.6. Nucleotide sequence accession number

A 1554-bp sequence including the *bla*_{KPC-2} gene corresponding to isolate ECL-13354 (ECL-P2) and KPN-13355 (KPN-P2) has been submitted to the GenBank nucleotide sequence database under accession nos. KR108242 and KR108243, respectively.

3. Results and discussion

Epidemiological data for the patients is shown in Table 1. The three patients had prolonged hospitalisation and serious underlying conditions. Patients 2 and 3 had had previous antibiotic treatment due to multiple nosocomial bacterial and fungal infections. Isolate recovery was 1 day and 7 days apart for Cases 1 and 2, respectively, and simultaneously in Case 3.

The susceptibility patterns of all isolates are shown in Table 2. Susceptibility to carbapenems was variable, from fully susceptible to resistant, whilst the isolates were highly resistant to expanded-spectrum β -lactams. Resistance to aminoglycosides and fluoroquinolones was also variable, whereas colistin and fosfomycin remained susceptible. All studied isolates harboured *bla*_{KPC-2}, and the only ESBL detected was *bla*_{PER-2} in isolates from Patient 3 (Table 2).

The *K. pneumoniae* isolates from Patients 1 and 2 were not clonally related (>6 band difference); however, the isolate recovered from Patient 1 was genetically related to *K. pneumoniae* ST258 (data not shown).

Transconjugants were obtained for the isolates from Patient 2 (TC-ECL-P2 and TC-KPN-P2) and Patient 3 (TC-CFR-P3 and TC-KOX-P3). All transconjugants were shown by PCR to contain *bla*_{KPC-2}. Carbapenem MICs of transconjugants obtained from Patient 2 were at least three times higher than those of the recipient cells (*E. coli* J53) (Table 2). Moreover, *bla*_{PER-2} was

Table 2 Susceptibility testing and interpretation, genes detected, plasmid profile and genetic context of *bla*_{KPC-2}.

| Patient | Isolate | MIC (μg/mL) [susceptibility interpretation] | | | | | | | | | | | | | | No. of plasmids and MW (kb) ^a | RT | Genetic element(s) |
|---------|--------------------|---|----------|-----------|----------|----------|----------|---------|----------|----------|----------|-----------|----------|----------|-----------------------------|--|----------|---|
| | | IPM | MER | ETP | FEP | CAZ | CTX | MIN | COL | FOS | TIG | CIP | AMK | GEN | | | | |
| 1 | ECO-P1 | 0.5 [S] | 0.5 [S] | 0.25 [S] | 2 [S] | 16 [I] | 4 [R] | 0.5 [S] | 0.25 [S] | 0.5 [S] | 0.25 [S] | 0.008 [S] | 64 [R] | 64 [R] | <i>bla</i> _{KPC-2} | <48.5, 70 | NT | Tn4401a |
| | KPN-P1 | 1 [S] | 1 [S] | 8 [S] | 16 [R] | >64 [R] | 16 [R] | 16 [R] | 0.5 [S] | 4 [S] | 1 [S] | ≥16 [R] | 64 [R] | 32 [R] | <i>bla</i> _{KPC-2} | <48.5, 58, 70, 211 | NT | |
| 2 | ECL-P2 | 4 [R] | 8 [R] | 16 [R] | ≥64 [R] | 64 [R] | ≥64 [R] | 16 [R] | 0.5 [S] | 2 [S] | 2 [I] | 8 [R] | 2 [S] | 0.5 [S] | <i>bla</i> _{KPC-2} | 65.5, 162 | L/M | ISKpn8- <i>bla</i> _{KPC-2} -ISKpn6-like |
| | TC-ECL-P2 | 4 [R] | 2 [I] | 1 [I] | 4 [S] | 4 [S] | 8 [R] | 1 [S] | 0.25 [S] | 0.5 [S] | 0.25 [S] | 0.015 [S] | 0.12 [S] | 0.12 [S] | <i>bla</i> _{KPC-2} | 65.5 | L/M | |
| | KPN-P2 | 2 [I] | 1 [S] | 4 [R] | 8 [I] | 32 [R] | 8 [R] | 8 [I] | 1 [S] | 8 [S] | 1 [S] | 0.25 [S] | 2 [S] | 0.5 [S] | <i>bla</i> _{KPC-2} | 70, 86 | L/M | |
| | TC-KPN-P2 | 4 [R] | 2 [I] | 1 [I] | 4 [I] | 4 [S] | 8 [R] | 8 [R] | 1 [S] | 0.25 [S] | 0.5 [S] | 0.008 [S] | 0.5 [S] | 0.12 [S] | <i>bla</i> _{KPC-2} | 70, 165 | L/M | |
| 3 | CFR-P3 | 0.5 [S] | 0.12 [S] | 0.12 [S] | 8 [I] | >64 [R] | 16 [R] | 32 [R] | 0.25 [S] | 0.25 [S] | 4 [R] | 8 [R] | 32 [I] | 8 [S] | <i>bla</i> _{KPC-2} | 70, 178–291 | A/C | ISKpn8- Δ <i>bla</i> _{TEM} - <i>bla</i> _{KPC-2} -ISKpn6-like |
| | TC-CFR-P3 | 0.25 [S] | 0.5 [S] | 0.125 [S] | 2 [S] | ≥128 [R] | 8 [R] | 1 [S] | 0.25 [S] | 0.5 [S] | 0.25 [S] | 0.015 [S] | 8 [S] | 2 [S] | <i>bla</i> _{KPC-2} | 70, 204 | A/C | |
| | KOX-P3 | 0.5 [S] | 0.12 [S] | 0.5 [S] | 8 [I] | >64 [R] | 16 [R] | 4 [S] | 0.25 [S] | 4 [S] | 0.5 [S] | ≥16 [R] | 16 [S] | ≥256 [R] | <i>bla</i> _{KPC-2} | 70, 118, 168 | A/C, L/M | |
| | TC-KOX-P3 | 0.5 [S] | 0.5 [S] | 0.25 [S] | 2 [S] | 64 [R] | 8 [R] | 1 [S] | 0.25 [S] | 0.5 [S] | 0.25 [S] | 1 [R] | 8 [S] | 4 [S] | <i>bla</i> _{KPC-2} | 76 | A/C, L/M | |
| | <i>E. coli</i> J53 | 0.25 [S] | 0.03 [S] | 0.008 [S] | 0.06 [S] | 0.25 [S] | 0.12 [S] | N/D | N/D | N/D | N/D | 0.015 [S] | 2 [S] | 0.25 [S] | N/A | N/A | N/A | N/A |

MIC, minimum inhibitory concentration; IPM, imipenem; MEM, meropenem; ETP, erTapentam; FEP, ceftazidime; CAZ, ceftazidime; CTX, cefotaxime; MIN, minocycline; COL, colistin; FOS, fosfomicin; TIG, tigecycline; CIP, ciprofloxacin; AMK, amikacin; GEN, gentamicin; RT, replicon typing; MW, molecular weight; R, resistant; I, intermediate; S, susceptible; N/D, not determined; N/A, not applicable.

^a Numbers in bold represent positive hybridisation bands with the *bla*_{KPC-2} probe.

co-transferred to the transconjugants. The different MICs observed for β -lactams of TC-ECL-P2 compared with the parental strain ECL-P2 were possibly due to the presence of the chromosomal AmpC of *E. cloacae*.

Table 2 shows the plasmid profile obtained after S1 nuclease digestion of total DNA and PBRT [12,13]. The six clinical isolates carried multiple plasmids ranging from 45 kb to 300 kb. In Case 1, the plasmids carrying *bla*_{KPC-2} in ECO-P1 and KPN-P1 were non-typeable by PBRT and had an approximate size of 70 kb, suggesting that both strains may harbour the same plasmid. In Case 2, *bla*_{KPC-2} was detected on L/M-type plasmids, slightly different in size (ca. 65 kb in ECL-P2 and TC-ECL-P2 and ca. 70 kb in KPN-P2 and TC-KPN-P2) (Table 2). In Case 3, CFR-P3 and KOX-P3 carried *bla*_{KPC-2} on a ca. 70 kb A/C-type plasmid (Table 2). In this particular case, comparison of the sizes of the plasmids in the clinical strains differed with those of the transconjugants. The S1-PFGE gel of TC-CFR-P3 showed two bands of 70 kb and 204 kb, respectively, and no specific hybridisation was observed, possibly due to the low-copy number plasmid, below the sensitivity of the technique. In TC-KOX-P3, the specific hybridisation band observed was of ca. 76 kb. The differences observed could be explained by plasmid rearrangements or transposition events (e.g. Case 3, 70 kb vs. 76 kb).

Analysis of the genetic elements surrounding *bla*_{KPC-2} using PCR and sequencing revealed the presence of three different genetic elements and insertion sequences. In Case 1, ECO-P1 and KPN-P1 harboured the already described transposon Tn4401a [14]. To the best of our knowledge, this is the first report of Tn4401 detected in *E. coli* in Argentina because this structure has always been associated with *K. pneumoniae* ST258 [10]. Isolates from Patient 2 and transconjugants shared a novel genetic context within the variable region between ISKpn-8 and *bla*_{KPC-2}, which we called Variant 3 as it differs from the Variant 2 published by Shen et al. [15] (GenBank FJ628167). These structures had a 114-bp deletion that included 24 bp of the inverted repeat right of the Tn3-like transposon (GenBank KM403446). Finally, isolates from Patient 3 harboured *bla*_{KPC-2} in the genetic context Variant 1a: ISKpn8- Δ *bla*_{TEM}-*bla*_{KPC-2}-ISKpn6-like (GenBank JN048639) already reported in Argentina [10].

In conclusion, this work documents three cases of in vivo horizontal transfer of *bla*_{KPC}. The results describe the variety of genetic elements harbouring *bla*_{KPC} that are circulating in Argentina as well as the first description of *bla*_{KPC}-producing *E. coli* associated with Tn4401a.

The broad dissemination of KPC-producing *K. pneumoniae* increases the likelihood of interspecies transfer of the antibiotic resistance determinants into highly fit clones of other Enterobacteriaceae. Therefore, early detection, characterisation and surveillance of these resistance elements are extremely important to avoid their dissemination and consequent treatment failures.

Funding

This work was supported by grants from Agencia Nacional de Promoción Científica y Tecnológica (ANPCYT) to SAG [BID-PICT-2008-0286 and PICT-2012-0145] and by the regular federal budget of the National Ministry of Health of Argentina.

Competing interests

None declared.

Ethical approval

Not required.

References

- [1] Ruiz-Garbajosa P, Curiao T, Tato M, Gijon D, Pintado V, Valverde A, et al. Multiclonal dispersal of KPC genes following the emergence of non-ST258 KPC-producing *Klebsiella pneumoniae* clones in Madrid, Spain. *J Antimicrob Chemother* 2013;68:2487–92.
- [2] Tsakris A, Voulgari E, Poulou A, Kimouli M, Pournaras S, Ranellou K, et al. In vivo acquisition of a plasmid-mediated *bla*_{KPC-2} gene among clonal isolates of *Serratia marcescens*. *J Clin Microbiol* 2010;48:2546–9.
- [3] Leao RS, Carvalho-Assef AP, Correal JC, Silva RV, Goldemberg DC, Asensi MD, et al. KPC-2 producing *Klebsiella pneumoniae* and *Escherichia coli* co-infection in a catheter-related infection. *Clin Microbiol Infect* 2011;17:380–2.
- [4] Gona F, Barbera F, Pasquariello AC, Grossi P, Gridelli B, Mezzatesta ML, et al. In vivo multiclonal transfer of *bla*_{KPC-3} from *Klebsiella pneumoniae* to *Escherichia coli* in surgery patients. *Clin Microbiol Infect* 2014;20:O633–5.
- [5] Mathers AJ, Cox HL, Kitchel B, Bonatti H, Brassinga AK, Carroll J, et al. Molecular dissection of an outbreak of carbapenem-resistant Enterobacteriaceae reveals intergenus KPC carbapenemase transmission through a promiscuous plasmid. *MBio* 2011;2:e00204–211.
- [6] Clinical and Laboratory Standards Institute. Methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically: approved standard. 9th ed. Wayne, PA: CLSI; 2012, Document M07-A9.
- [7] Clinical and Laboratory Standards Institute. Performance standards for antimicrobial susceptibility testing: twenty-fourth informational supplement. Wayne, PA: CLSI; 2015, Document M100-S24.
- [8] Pasteran F, Mendez T, Guerriero L, Rapoport M, Corso A. Sensitive screening tests for suspected class A carbapenemase production in species of Enterobacteriaceae. *J Clin Microbiol* 2009;47:1631–9.
- [9] Lee K, Lim YS, Yong D, Yum JH, Chong Y. Evaluation of the Hodge test and the imipenem–EDTA double-disk synergy test for differentiating metallo- β -lactamase-producing isolates of *Pseudomonas* spp. and *Acinetobacter* spp. *J Clin Microbiol* 2003;41:4623–9.
- [10] Gomez SA, Pasteran FG, Faccone D, Tijet N, Rapoport M, Lucero C, et al. Clonal dissemination of *Klebsiella pneumoniae* ST258 harbouring KPC-2 in Argentina. *Clin Microbiol Infect* 2011;17:1520–4.
- [11] Tenover FC, Arbeit RD, Goering RV, Mickelsen PA, Murray BE, Persing DH, et al. Interpreting chromosomal DNA restriction patterns produced by pulsed-field gel electrophoresis: criteria for bacterial strain typing. *J Clin Microbiol* 1995;33:2233–9.
- [12] Carattoli A, Bertini A, Villa L, Falbo V, Hopkins KL, Threlfall EJ. Identification of plasmids by PCR-based replicon typing. *J Microbiol Methods* 2005;63:219–28.
- [13] Barton BM, Harding GP, Zuccarelli AJ. A general method for detecting and sizing large plasmids. *Anal Biochem* 1995;226:235–40.
- [14] Naas T, Cuzon G, Villegas MV, Lartigue MF, Quinn JP, Nordmann P. Genetic structures at the origin of acquisition of the β -lactamase *bla*_{KPC} gene. *Antimicrob Agents Chemother* 2008;52:1257–63.
- [15] Shen P, Wei Z, Jiang Y, Du X, Ji S, Yu Y, et al. Novel genetic environment of the carbapenem-hydrolyzing β -lactamase KPC-2 among Enterobacteriaceae in China. *Antimicrob Agents Chemother* 2009;53:4333–8.